



Lipophilic compounds in femoral secretions of males and females of the El Hierro giant lizard *Gallotia simonyi* (Lacertidae)



José Martín ^{a,*}, Albert Martínez-Silvestre ^b, Pilar López ^a, Alejandro Ibáñez ^a, Miguel Ángel Rodríguez-Domínguez ^c, Isabel Verdaguer ^b

^a Departamento de Ecología Evolutiva, Museo Nacional de Ciencias Naturales, CSIC, José Gutiérrez Abascal 2, 28006 Madrid, Spain

^b CRARC, Catalonian Reptile and Amphibian Recovery Center, 08783 Masquefa, Barcelona, Spain

^c Cabildo Insular de El Hierro, Centro de Reproducción e Investigación del lagarto gigante de El Hierro, El Hierro, Canary Islands, Spain

ARTICLE INFO

Article history:

Received 27 February 2015

Received in revised form 25 June 2015

Accepted 26 June 2015

Available online xxx

Keywords:

Lizards

Gallotia simonyi

Femoral secretions

Intersexual variations

Canary Islands

ABSTRACT

Many lizards use femoral gland secretions in reliable intraspecific communication. Based on mass spectra, obtained by GC–MS, we found 57 lipophilic compounds in femoral secretions of males and females of El Hierro giant lizards, *Gallotia simonyi* (fam. Lacertidae). Compounds included steroids (mainly cholesterol) and fatty acids ranging between *n*-C₁₆ and *n*-C₂₂ (mainly hexadecanoic and octadecanoic acids), followed by aldehydes, alcohols, ketones, squalene and waxy esters. There were important intersexual differences in the presence, abundance and number of compounds (more numerous in males). Males had higher proportions of the most odoriferous compounds (fatty acids and aldehydes), while females had higher proportions of more stable compounds (steroids, waxy alcohols, waxy esters and terpenoids). This suggests sexual differences in function of femoral secretions. In addition, some compounds could reflect the physiological state, allowing monitoring health of lizards from secretions samples, which is especially important given the critical conservation status of this lizard.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

Chemical signals are important for intraspecific communication of many animals (Müller-Schwarze, 2006; Wyatt, 2014). For example, lizards produce femoral or precloacal gland secretions with chemical compounds that are used in communication and social organization and may have consequences for reproduction (Mason, 1992; Alberts, 1993; Mason and Parker, 2010; Martín and López, 2011, 2014). In many lizard species, males use femoral secretions to scent-mark territories, and lipophilic compounds in these scent-marks may provide information on the characteristics, dominance status and quality of the territory owner to other males (López and Martín, 2002, 2011; Labra, 2006; Carazo et al., 2007; Martín and López, 2007; Martín et al., 2007b; Khannoob et al., 2011b) and to females (Martín and López, 2000, 2006a,b; 2012; Olsson et al., 2003; López and Martín, 2005a, 2012). This information is reliable because the amount of some compounds that can be allocated to secretions is dependent on conflicting physiological requirements, such as the maintenance of the immune system,

* Corresponding author.

E-mail address: Jose.Martin@mncn.csic.es (J. Martín).

and it also depends on the diet and the endocrine and health state of males ([Alberts et al., 1992a; López et al., 2006, 2009; Martín et al., 2007a; Kopena et al., 2011, 2014; Heathcote et al., 2014](#); see review in [Martín and López, 2015](#)). Therefore, to understand the role of chemical signals in reproduction and sexual selection of a lizard species, we need first to characterize the compounds secreted by the femoral glands and their variation.

Femoral secretions of lizards are composed of proteins and many lipophilic compounds, such as steroids, fatty acids, alcohols, etc (reviewed in [Weldon et al., 2008; Martín and López, 2011, 2014](#)). However, the specific chemical compounds found in each species and their relative proportions are only known for a few species, which are mainly lacertids ([López and Martín, 2005b,c, 2006; 2009; Martín and López, 2006c,d; 2010; Gabiro et al., 2008, 2010, 2012b; Kopena et al., 2009; Khanno et al., 2011a; Martín et al., 2013a](#)) and several South American tropidurids of the genus *Liolaemus* ([Escobar et al., 2001, 2003](#)), although there is also information on an African cordylid ([Louw et al., 2007](#)), an American teiid ([Martín et al., 2011](#)), two iguanids ([Weldon et al., 1990; Alberts et al., 1992a,b; Martín et al., 2013b](#)), three agamids ([Chauhan, 1986; Martín et al., 2012, 2013c](#)) and a few gekkonids ([Chauhan, 1986; Khanno, 2012](#)). These studies show consistent interspecific variation in composition ([Weldon et al., 2008; Martín and López, 2011, 2014](#)), which probably reflects phylogenetic differences *per se*, although environmental conditions might also contribute to explain why some compounds have been selected in a given species ([Martín and López, 2006c, 2013](#)). Also, because allocation of compounds to secretions may depend on the physiological state of an individual (e.g. [Martín et al., 2007a; Martín and López, 2015](#)), examining interindividual variation in chemical compounds may also be useful in other applied contexts, such as in monitoring of populations.

The El Hierro Giant Lizard (*Gallotia simonyi*), is a large sized lacertid species which was believed to be extinct in the 1930s but that was rediscovered in 1974 ([Böhme and Bings, 1975; Barahona et al., 2000](#)). It is included in the genus *Gallotia*, endemic from the Canary Islands (Spain) ([Arnold, 1973](#)). This species is restricted to El Hierro Island (Canary Islands) where it is now confined to a few small areas of a high cliff ("Risco de Tibataje") with sparse vegetation, although new populations have been recently reintroduced in other areas. The population in the wild is now about 300 animals (including re-introduced populations) ([Rodríguez-Domínguez et al., 2014](#)). The conservation status of *G. simonyi* is classified as "critical" by the International Union for Conservation of Nature (IUCN) guidelines ([Pleguezuelos et al., 2002](#)). Thus, this lizard is strictly protected by national and international laws and is object of a recovery plan, that includes a captivity breeding program and a regular monitoring of physiological health state of lizards.

Several articles have examined physiological and clinical aspects of *G. simonyi*, such as hematology ([Martínez-Silvestre et al., 2004a, 2005](#)), microbiology ([Martínez-Silvestre et al., 2004b; Herrera-León et al., 2005](#)), reproduction ([Rodríguez-Domínguez and Molina-Borja, 1998](#)), diseases ([Martínez-Silvestre and Orós, 2002](#)) and parasitology ([Roca et al., 1999](#)), as well as some ethological aspects ([Cejudo et al., 1997](#)). To date, nothing is known on the role of chemical ecology in social behavior and sexual selection of Canarian lizards of the genus *Gallotia*, or of *G. simonyi* in particular.

We report here the results of an analysis by gas chromatography-mass spectrometry (GC-MS) of the lipophilic fraction of femoral secretions of adult males and females of El Hierro giant lizards, *G. simonyi*. We specifically describe and compare between sexes the composition and relative proportions of compounds in femoral secretions of lizards.

2. Material and methods

2.1. Study animals

The lizards studied here (22 males and 9 females) came from the Recovery Centre for the Giant Lizard of El Hierro (El Hierro, Canary Islands, 14° 00' W; 27° 45' N), located in the same area where natural populations of this lizard can be found. These animals are part of captive breeding programme, which originally used 12 wild animals captured between 20 and 30 years ago. New introduced populations have been established using lizards from this captive stock. The existence of relatively few adult examples for the breeding programme is due to the scarcity of wild adults compatible with a maximum genetic variability that would allow the viability of the species ([Arano et al., 1997](#)).

In this recovery centre, lizards were kept in pairs, in large outdoor seminatural enclosures under the same temperature and light conditions that they experienced in the wild. During the study period, temperature range was between 23 and 35 °C (night-day) and photoperiod was 14 h light and 10 h dark. Lizards were fed regularly (three times at week), on an omnivorous diet of fruit, vegetables and endemic plants, supplemented with crickets (*Grillus bimaculatus*) and giant mealworms (*Zophobas morio*). This diet resembles the natural diet of lizards in the wild ([Pérez-Mellado et al., 1999](#)). The lizards were all eating, drinking and behaving normally at the time of the study.

Lizards sampled were aged between four and twelve years. The study was performed in summer 2014, and most animals still maintained abundant active femoral secretions. However, females had inactive ovaries, and abdominal palpation of each one confirmed that they had no oviductal eggs.

2.2. Chemical analyses of femoral gland secretions

Samples of femoral secretions were collected in an active period of lizards (10.00–13.00 h) after they had finished their morning basking periods. We extracted secretions from femoral glands of lizards by gently pressing with forceps around the femoral pores. The secretion was directly collected into glass inserts contained in glass vials, which were later closed with Teflon-lined stoppers. Vials were stored at –20 °C until analyses. We also made the same procedure but without collecting

secretion, to obtain blank control vials that were used to exclude contaminants from the handling procedure or from the environment, and for examining potential impurities in the solvent or laboratory equipment.

For chemical analyses of samples, we used a Finnigan-ThermoQuest Trace 2000 gas chromatograph (GC) fitted with a poly(5% diphenyl/95% dimethylsiloxane) column (Supelco, Equity-5, 30 m length × 0.25 mm ID, 0.25-μm film thickness) and a Finnigan-ThermoQuest Trace mass spectrometer (MS) as detector. We injected 2 μl of each sample dissolved in *n*-hexane (Sigma, capillary GC grade, 99.9% purity), using the splitless mode and helium as the carrier gas at 30 cm/s. The temperature of the injector was set at 300 °C. The oven temperature program started at 50 °C, was maintained isothermal for 3 min, then increased to 300 °C at a rate of 5 °C/min, and then isothermal (300 °C) for 15 min. Ionization by electron impact (70 eV) was carried out at 280 °C. Mass spectral fragments below *m/z* = 40 were not recorded. Impurities identified in the control vial samples are not reported. We made initial identification of secretions components by comparison of mass spectra in the NIST/EPA/NIH (NIST 02) computerized mass spectral library. Identifications were confirmed by comparison of spectra and retention times with those of authentic standards (from Sigma–Aldrich Chemical Co) when these were available.

2.3. Data analyses

We determined the relative amount of each compound in the secretions as the percent of the total ion current (TIC). To correct the problem of nonindependence of proportions, we transformed data using the compositional analysis ($\ln(\text{proportion})/(1 - \text{proportion})$) (Aebischer et al., 1993). Then, to test whether the chemical profiles differed between males and females, we used a single factor permutational multivariate analysis of variance test (PERMANOVA) using 999 permutations, (Anderson, 2001; McArdle and Anderson, 2001) based on a resemblance matrix produced by calculating Euclidean distances between every pair of individual samples. To calculate whether a chemical profile could be assigned to a sex based on its characteristics, we used a canonical analysis of principal coordinates (CAP, Anderson and Willis, 2003) and a permutation test. We used for these analyses the software PRIMER V6.1.13 (Clarke and Gorley, 2006) with the PERMANOVA V1.0.3 add-on package (Anderson et al., 2008).

3. Results

We found 57 lipophilic compounds in femoral gland secretions of male *G. simonyi* (Table 1). The main components were 22 steroids (43.1% of TIC) and nine carboxylic acids or their esters ranged between *n*-C₁₆ and *n*-C₂₂ (41.5%), but we also found ten aldehydes (6.9%), six alcohols between C₁₅ and C₂₂ (5.9%), squalene (1.3%), three ketones between C₁₇ and C₂₁ (0.7%) and five waxy esters (0.6%). On average, the five most abundant chemicals in males were cholesterol (24.9% of TIC) and octadecanoic acid (22.1%), followed by eicosanoic acid (6.7%), sitosterol (6.1%) and hexadecanol (4.9%).

In contrast, in females we only found 27 lipophilic compounds (Table 1). The main components were six steroids (50.3% of TIC), but we also found six alcohols between C₁₅ and C₂₄ (22.3%), squalene and other unidentified terpenoid (12.4%), six waxy esters (9.7%), four aldehydes (3.5%), two esters of carboxylic acids (1.4%), and one ketone (0.4%). On average, the most abundant chemicals in females were cholesterol (36.9% of TIC) and tetracosanol (14.8%), followed by squalene (12.3%), cholest-4-en-3-one (5.4%) and hexadecanol (4.7%).

Only 21 of the 63 identified compounds (33.3%) were shared by both sexes, which comprised 43.1% of TIC areas of males. Males had 36 exclusive compounds (63.2% of compounds) that comprised 56.9% of the TIC areas of males, while females had only six exclusive compounds (22.2%) that comprised 22.3% of their TIC areas. There were significant differences in the multivariate comparison of the chemical profiles of femoral secretions between sexes (PERMANOVA based on the resemblance matrix, pseudo $F_{1,28} = 43.30$, $p = 0.001$), and 100% of the chemical profiles could be classified into the correct sex (CAP analysis using leave-one-out cross-validation, $\delta_1^2 = 0.91$, $p = 0.001$, $m = 2$ axes).

Comparing the relative abundance (% of TIC) of the different type of compounds between sexes, males had significantly higher proportions of fatty acids (GLM, $F_{1,28} = 64.71$, $p < 0.0001$) and aldehydes ($F_{1,28} = 6.05$, $p = 0.02$) than females. In contrast, females had significantly more steroids ($F_{1,28} = 6.50$, $p = 0.017$), alcohols ($F_{1,28} = 31.89$, $p < 0.0001$), waxy esters ($F_{1,28} = 105.00$, $p < 0.0001$) and terpenoids ($F_{1,28} = 25.16$, $p < 0.0001$) than males.

4. Discussion

The lipophilic fraction of femoral secretions of *G. simonyi* is formed mainly by steroids and fatty acids, with cholesterol being the main compound, as it is usual in other lacertid lizards (Weldon et al., 2008). Also, there were minor amounts of other types of compounds, such as alcohols, aldehydes, terpenoids, etc that are also usually found in other related lizards. In addition, there were important intersexual differences in presence and relative proportions of compounds.

With respect to steroids, the large amount of cholesterol, as in other lizards, may form an unreactive apolar matrix to stabilize secretions and protect other compounds with a signaling function (Escobar et al., 2003). This would be especially important in the hot and dry habitats with high evaporation rates, where *G. simonyi* lives (Alberts, 1992; Gabirov et al., 2012a). Other steroids with a relative high abundance are sitosterol and campesterol, steroids of vegetal origin (i.e., phytosterols) that have to be obtained from the diet and diverted from their potentially beneficial physiological functions (Choudhary and Tran, 2011; Genser et al., 2012) to be allocated to secretions. Thus, the presence and relative amounts of these compounds in

Table 1Lipophilic compounds found in femoral secretions of male and female El Hierro Giant lizards, *Gallotia simonyi*.

RT (min)	Compound	Males (n = 22) mean ± SE	Females (n = 9) mean ± SE
26.0	Tetradecanal	0.05 ± 0.03	—
27.7	Hexadecenal	2.06 ± 0.44	0.75 ± 0.04
28.2	Pentadecanal	0.21 ± 0.04	0.18 ± 0.18
29.9	Pentadecanol	0.09 ± 0.02	0.95 ± 0.62
30.4	Hexadecanal	1.42 ± 0.46	0.81 ± 0.24
30.9	Unidentified terpenoid	—	0.09 ± 0.09
31.4	Hexadecenol	0.01 ± 0.01	—
31.6	Hexadecenal	0.01 ± 0.01	—
31.9	Hexadecanol	4.95 ± 0.81	4.74 ± 0.84
32.1	2-Heptadecanone	0.22 ± 0.06	—
32.4	Heptadecanal	0.02 ± 0.02	—
32.6	14-Methyl-hexadecanoic acid, methyl ester	0.03 – 0.03	0.17 ± 0.17
33.1	9-Hexadecenoic acid	0.04 ± 0.03	—
33.3	Hexadecanoic acid	4.28 ± 0.27	—
33.9	Octadecenal	0.10 – 0.06	—
34.4	Octadecanal	2.38 ± 0.74	1.75 ± 0.41
34.4	1-Methylethyl ester-hexadecanoic acid	0.11 ± 0.09	1.28 ± 0.52
35.7	Octadecanol	0.58 ± 0.11	0.45 ± 0.45
35.9	2-Nonadecanone	0.40 ± 0.12	0.40 ± 0.40
36.0	5-Dodecyldihydro-2(3H)-furanone	0.14 ± 0.04	—
36.3	Nonadecanal	0.04 ± 0.04	—
36.7	9-Octadecenoic acid	1.48 ± 0.30	—
37.1	Octadecanoic acid	22.09 ± 1.77	—
37.9	6-Octadecenoic acid	4.37 ± 1.60	—
38.0	Eicosanal	0.62 ± 0.34	—
39.2	Eicosanol	0.14 ± 0.11	0.27 ± 0.27
39.4	2-Heneicosanone	0.05 ± 0.05	—
40.5	Eicosanoic acid	6.68 ± 2.87	—
41.5	Docosenoic acid	2.40 ± 1.69	—
42.6	Docosanol	0.09 ± 0.09	1.11 ± 1.11
47.3	Squalene	1.29 ± 0.79	12.29 ± 8.92
47.9	Cholesta-2,4-diene	0.24 ± 0.17	—
48.1	Cholesta-4,6-dien-3-ol	0.94 ± 0.22	—
48.2	Tetracosanol	—	14.78 ± 12.48
48.4	Cholesta-3,5-diene	0.70 ± 0.13	—
48.7	Unid.Steroid (197,251,355,364,387,400,414)	0.02 ± 0.02	—
49.6	Cholesta-4,6-dien-3-ol, methyl derivative?	0.23 ± 0.18	—
50.0	Unid.Steroid (215,253,376,393)	0.02 ± 0.02	—
50.6	Unid.Steroid (197,251,371,382,390,396)	0.13 ± 0.06	—
50.8	Cholesta-4,6-dien-3-ol, ethyl derivative?	0.51 ± 0.24	—
51.0	Cholesta-3,5-diene, ethyl derivative?	0.33 ± 0.13	—
51.4	Cholesterol	24.94 ± 2.96	36.90 ± 3.72
51.5	Cholestanol	—	3.64 ± 0.06
51.9	Cholestan-3-one, methyl derivative?	—	1.44 ± 0.46
52.0	Cholesta-5,24-dien-3-ol (=Desmosterol)	1.60 ± 0.20	—
52.8	Campesterol	2.41 ± 0.22	1.64 ± 0.08
53.0	Cholest-4-en-3-one	0.76 ± 0.22	5.41 ± 1.61
53.2	Stigmasterol	1.31 ± 0.18	—
53.3	Ergost-22-en-3-ol	0.02 ± 0.02	—
53.4	Cholesta-4,6-dien-3-one	0.74 ± 0.21	1.28 ± 1.28
53.7	Unid.Steroid (211,253,271,351,377,411)	0.06 ± 0.06	—
53.8	Hexadecyl 9-hexadecenoate	0.15 ± 0.15	1.58 ± 0.51
53.9	Sitosterol	6.11 ± 0.92	—
54.0	Hexadecyl hexadecanoate	—	2.33 ± 0.65
54.1	Stigmasterol	1.30 ± 0.21	—
54.4	9-Octadecenyl 9-hexadecenoate	0.05 ± 0.05	—
54.4	20-Methyl-pregn-20-en-3-ol	0.09 ± 0.06	—
54.8	Ergostenol, methyl derivative?	0.41 ± 0.11	—
54.9	Octadecyl 9-hexadecenoate	0.13 ± 0.13	0.92 ± 0.92
55.3	Nonadecyl 9-hexadecenoate	—	1.47 ± 0.71
56.3	Cholesta-4,6-dien-3-one, derivative?	0.21 ± 0.11	—
56.9	Eicosyl 9-hexadecenoate	0.16 ± 0.11	1.34 ± 1.34
57.2	Octadecyl hexadecanoate	0.08 ± 0.07	2.02 ± 0.78

The relative amount of each component was determined as the percent of the total ion current (TIC) and reported as the average ($\pm 1\text{SE}$). Characteristic ions (*m/z*) are reported for unidentified compounds. RT: Retention time.

secretions might be directly related to the characteristics, and may be quality, of the diet of each individual lizard (Martín and López, 2015).

The fatty acids that appear in secretions are those usually found in other lizards (Weldon et al., 2008; Martín and López, 2014). However, the high relative proportion of octadecanoic acid in males is notorious, when comparing it with most of other lacertid lizard species, where hexadecanoic acid is more abundant (e.g. López and Martín, 2005b; Martín and López, 2006c). Nevertheless, this fatty acid is also the most abundant one in secretions of the lacertid lizard *Psammodromus algirus* (Martín and López, 2006d). *Psammodromus* is considered the genus most phylogenetically related to the genus *Gallotia*, both forming the subfamily Gallotiinae, which shows a great genetic divergence from the rest of lacertids (Arnold, 1973; Mayer and Pavlicev, 2007). This suggests that phylogenetic differences *per se* between lizard species might account for a high percentage of the observed interspecific variability in femoral secretion composition. Nevertheless, octadecanoic acid is not so especially abundant in other species of the genus *Gallotia* from others of the Canary Islands (J. Martín unpubl. data), suggesting that the environment also modulates the compounds found in lizards' secretions.

Interestingly, in the related *P. algirus*, individual males with lower blood parasite loads and greater immune responses (i.e. with a better health state) have femoral secretions with lower proportions of two carboxylic acids (octadecanoic and eicosanoic acids) and higher proportions of their correspondent alcohols (octadecanol and eicosanol) (Martín et al., 2007a). Moreover, female *P. algirus* discriminate these compounds and respond with higher chemosensory rates to secretions of healthier males, suggesting that the scents of these males are more attractive (Martín et al., 2007a). A similar, yet unexplored, signaling function of these compounds might occur in *G. simonyi*, which prompts for the design of future experiments to unravel the function of femoral secretions in this lizard. Also, data from femoral secretions composition might complement the already-known physiological parameters and be useful in future studies to estimate in an innocuous way the health state of lizards in both wild and captive populations of this highly threatened species.

In most species of lizards with femoral glands, females also possess femoral pores, but they are usually much smaller and with very little or without apparent secretion (Mason, 1992). Femoral secretion of female *G. simonyi* is also less abundant than that of males, but the large body size of this lizard allowed us to collect and analyze it. The chemical analysis showed important differences in composition, with females having a lower number of compounds than males and with only a few of them being exclusive of females. But intersexual differences did not arise just because the lack of some compounds in females, but also because some of the main shared compounds and the different major types of compounds had very different proportions. In contrast, in the other lacertid, *Acanthodactylus boskianus*, where secretions of females have been analyzed, the same compounds are found in both sexes, although there are some small differences in proportions of compounds (Khamnoon et al., 2011a). Nevertheless, it is possible that the lack of many compounds in our sample of female *G. simonyi* might be explained by seasonal differences in the production of femoral secretions of females, as it has been observed in iguanian lizards (Alberts, 1990; Kelso and Martins, 2008), although the specific secreted compounds were not analyzed in these species.

Intersexual differences in *G. simonyi* allowed predicting statistically the sex of an individual based on its chemical profile, and probably also allow lizards to easily discriminate by chemosensory cues the sex of conspecifics. The occurrence of this chemosensory ability has not been experimentally tested in this species, although it is expected based on the ability of *G. simonyi* to discriminate chemical cues of its prey (Cooper and Pérez-Mellado, 2001) and on the well known sex discriminatory abilities of most lacertid lizard species (Mason and Parker, 2010; Martín and López, 2011). Sex discrimination based on chemical cues could be especially important in intraspecific relationships of this species where both sexes are similar in coloration, being both black with some yellow ocelli, only differing in average body size and relative head size (Rodríguez-Domínguez et al., 1998).

Among the most important intersexual differences, females had high proportions of compounds that would confer a high stability to the secretions, notably a high proportion of a waxy alcohol (i.e. tetracosanol), that occurred only in females, and more cholesterol, squalene and waxy esters than males. In contrast, females lacked or had lower proportions than males of the more odoriferous compounds, such as fatty acids and aldehydes, which could be used in intraspecific communication at short-range. This suggests an entirely different role of femoral secretions in males and females of this lizard that merits further studies.

The results of this study suggest that if compounds in secretions of this lizard could reflect their physiological health state, future studies should establish which reliable relationships could be inferred from the chemical study of femoral secretions. This could allow monitoring the health state of lizards from a simple sample of their femoral secretions, avoiding other more invasive techniques such as blood sampling, which is especially important given the highly threatened status of *G. simonyi*.

Acknowledgements

We thank Elena Fernández and Luis Cuadra for technical assistance with chemical analyses. Financial support was provided by the project MCIIN-CGL2011-24150/BOS. Lizards were handled and sampled thanks to special permits from Cabildo de El Hierro.

References

- Aebischer, N.J., Robertson, P.A., Kenward, R.E., 1993. Compositional analysis of habitat use from animal radio-tracking data. *Ecology* 74, 1313–1325.
- Alberts, A.C., 1990. Chemical properties of femoral gland secretions in the desert iguana, *Dipsosaurus dorsalis*. *J. Chem. Ecol.* 16, 13–25.

- Alberts, A.C., 1992. Constraints on the design of chemical communication systems in terrestrial vertebrates. *Am. Nat.* 139, S62–S89.
- Alberts, A.C., 1993. Chemical and behavioral studies of femoral gland secretions in iguanid lizards. *Brain Behav. Evol.* 41, 255–260.
- Alberts, A.C., Pratt, N.C., Phillips, J.A., 1992a. Seasonal productivity of lizard femoral glands: relationship to social dominance and androgen levels. *Physiol. Behav.* 51, 729–733.
- Alberts, A.C., Sharp, T.R., Werner, D.I., Weldon, P.J., 1992b. Seasonal variation of lipids in femoral gland secretions of male green iguanas (*Iguana iguana*). *J. Chem. Ecol.* 18, 703–712.
- Anderson, M.J., 2001. A new method for non-parametric multivariate analysis of variance. *Austral Ecol.* 26, 32–46.
- Anderson, M.J., Willis, T.J., 2003. Canonical analysis of principal coordinates: a useful method of constrained ordination for ecology. *Ecology* 84, 511–525.
- Anderson, M.J., Gorley, R.N., Clarke, K.R., 2008. PERMANOVA β for PRIMER: Guide to Software and Statistical Methods. PRIMER-E Ltd, Plymouth, UK.
- Arano, B., Herrero, P., Astudillo, G., 1997. Gestión genética de una especie amenazada: monitorización de la variación genética y cría en cautividad del Lagarto Gigante de El Hierro, *Gallotia simonyi* (Sauria: Lacertidae). In: López Jurado, L.F., Mateo, J.A. (Eds.), El Lagarto Gigante de El Hierro. Bases para su Conservación. Monografías de Herpetología. Asociación Herpetológica Española, Madrid, Spain, pp. 85–100.
- Arnold, E.N., 1973. Relationships of the palearctic lizards assigned to the genera *Lacerta*, *Algyroides* and *Psammodromus* (Reptilia: Lacertidae). *Bull. Br. Mus. Nat. Hist. Zool.* 25, 1–366.
- Barahona, F., Evans, S.E., Mateo, J.A., García-Márquez, M., López-Jurado, L.F., 2000. Endemism, gigantism and extinction in island lizards: the genus *Gallotia* on the Canary Islands. *J. Zool.* 250, 373–388.
- Böhme, W., Bings, W., 1975. Zur frage des überlebens von *Lacerta s. simonyi* Steindachner (Sauria: Lacertidae). *Salamandra* 11, 39–46.
- Carazao, P., Font, E., Desfilis, E., 2007. Chemosensory assessment of rival competitive ability and scent mark function in a lizard (*Podarcis hispanica*). *Anim. Behav.* 74, 895–902.
- Cejudo, D., Marquez, R., García-Marquez, M., Bowker, R.G., 1997. Catálogo comportamental de *Gallotia simonyi*, el lagarto gigante de El Hierro (Islas Canarias). *Rev. Esp. Herp.* 11, 7–17.
- Chauhan, N.B., 1986. A preliminary report on the lipid components of pre-anal gland secretion of lizards. *Hemidactylus flaviviridis* and *Uromastix hardwickii*. *J. Anim. Morph. Phys.* 33, 73–76.
- Choudhary, S.P., Tran, L.S., 2011. Phytosterols: perspectives in human nutrition and clinical therapy. *Curr. Med. Chem.* 18, 4557–4567.
- Clarke, K.R., Gorley, R.N., 2006. PRIMER V6: User Manual/Tutorial. PRIMER-E Ltd, Plymouth, UK.
- Cooper Jr., W.E., Pérez-Mellado, V., 2001. Omnivorous lacertid lizards (*Gallotia*) from El Hierro, Canary Islands, can identify prey and plant food using only chemical cues. *Can. J. Zool.* 79, 881–887.
- Escobar, C.A., Labra, A., Niemeyer, H.M., 2001. Chemical composition of precloacal secretions of *Liolaemus* lizards. *J. Chem. Ecol.* 27, 1677–1690.
- Escobar, C.M., Escobar, C.A., Labra, A., Niemeyer, H.M., 2003. Chemical composition of precloacal secretions of two *Liolaemus fabiani* populations: are they different? *J. Chem. Ecol.* 29, 629–638.
- Gabiro, M., López, P., Martín, J., de Fraipont, M., Heulin, B., Sinervo, B., Cloibert, J., 2008. Chemical composition of femoral secretions of oviparous and viviparous types of male common lizards *Lacerta vivipara*. *Biochem. Syst. Ecol.* 36, 539–544.
- Gabiro, M., Castilla, A.M., López, P., Martín, J., 2010. Differences in chemical signals may explain species recognition between an island lizard, *Podarcis atrata*, and related mainland lizards, *P. hispanica*. *Biochem. Syst. Ecol.* 38, 521–528.
- Gabiro, M., López, P., Martín, J., 2012a. Interpopulational variation in chemosensory responses to selected steroids from femoral secretions of male lizards, *Podarcis hispanica*, mirrors population differences in chemical signals. *Chemoecology* 22, 65–73.
- Gabiro, M., López, P., Martín, J., 2012b. Differences in chemical sexual signals may promote reproductive isolation and cryptic speciation between Iberian wall lizard populations. *Int. J. Evol. Biol.* 2012, 1–13, article ID 698520.
- Genser, B., Silbernagel, G., De Backer, G., Bruckert, E., Carmen, R., Chapman, M.J., Deanfield, J., Descamps, O.S., Rietzschel, E.R., Dias, K.C., März, W., 2012. Plant sterols and cardiovascular disease: a systematic review and meta-analysis. *Eur. Heart J.* 33, 444–451.
- Heathcote, R.J.P., Bell, E., d'Errorre, P., While, G.M., Uller, T., 2014. The scent of sun worship: basking experience alters scent mark composition in male lizards. *Behav. Ecol. Sociobiol.* 68, 861–870.
- Herrera-León, S., Saco, M., Martínez-Silvestre, A., Silveira, L., Echeita, A., Usera, M.A., 2005. Molecular characterization of a new serovar of *Salmonella bongori* 13:22:Z39:- isolated from a lizard. *Res. Microbiol.* 156, 597–602.
- Kelso, E.C., Martins, E.P., 2008. The effects of two courtship display components on female reproductive behaviour and physiology in the sagebrush lizard. *Anim. Behav.* 75, 639–646.
- Khanno, E.R., 2012. Secretions of pre-anal glands of house-dwelling geckos (Family: Gekkonidae) contain monoglycerides and 1,3-alkanediol. A comparative chemical ecology study. *Biochem. Syst. Ecol.* 44, 341–346.
- Khanno, E.R., Flachsbarth, B., El-Gendy, A., Mazik, J., Hardege, J.D., Schulz, S., 2011a. New compounds, sexual differences, and age-related variations in the femoral gland secretions of the lacertid lizard *Acanthodactylus boskianus*. *Biochem. Syst. Ecol.* 39, 95–101.
- Khanno, E.R., El-Gendy, A., Hardege, J.D., 2011b. Scent marking pheromones in lizards: cholesterol and long chain alcohols elicit avoidance and aggression in male *Acanthodactylus boskianus* (Squamata: Lacertidae). *Chemoecology* 21, 143–149.
- Kopena, R., López, P., Martín, J., 2009. Lipophilic compounds from the femoral gland secretions of male Hungarian green lizards, *Lacerta viridis*. *Z. Naturforsch. C* 64, 434–440.
- Kopena, R., Martín, J., López, P., Herczeg, G., 2011. Vitamin E supplementation increases the attractiveness of males' scent for female European green lizards. *PLoS One* 6 (4), e19410.
- Kopena, R., López, P., Martín, J., 2014. Relative contribution of dietary carotenoids and vitamin E to visual and chemical sexual signals of male Iberian Green Lizards: an experimental test. *Behav. Ecol. Sociobiol.* 68, 571–581.
- Labra, A., 2006. Chemoreception and the assessment of fighting abilities in the lizard *Liolaemus monticola*. *Ethology* 112, 993–999.
- López, P., Martín, J., 2002. Chemical rival recognition decreases aggression levels in male Iberian wall lizards, *Podarcis hispanica*. *Behav. Ecol. Sociobiol.* 51, 461–465.
- López, P., Martín, J., 2005a. Female Iberian wall lizards prefer male scents that signal a better cell-mediated immune response. *Biol. Lett.* 1, 404–406.
- López, P., Martín, J., 2005b. Chemical compounds from femoral gland secretions of male Iberian rock lizards, *Lacerta monticola cyreni*. *Z. Naturforsch. C* 60, 632–636.
- López, P., Martín, J., 2005c. Age related differences in lipophilic compounds found in femoral gland secretions of male Spiny-footed lizards, *Acanthodactylus erythrurus*. *Z. Naturforsch. C* 60, 915–920.
- López, P., Martín, J., 2006. Lipids in the femoral gland secretions of male Schreiber's green lizards, *Lacerta schreiberi*. *Z. Naturforsch. C* 61, 763–768.
- López, P., Martín, J., 2009. Lipids in femoral gland secretions of male lizards, *Psammodromus hispanicus*. *Biochem. Syst. Ecol.* 37, 304–307.
- López, P., Martín, J., 2011. Male Iberian rock lizards may reduce the costs of fighting by scent-matching of the resource holders. *Behav. Ecol. Sociobiol.* 65, 1891–1898.
- López, P., Martín, J., 2012. Chemosensory exploration of male scent by female rock lizards result from multiple chemical signals of males. *Chem. Sens.* 37, 47–54.
- López, P., Amo, L., Martín, J., 2006. Reliable signaling by chemical cues of male traits and health state in male lizards, *Lacerta monticola*. *J. Chem. Ecol.* 32, 473–488.
- López, P., Gabiro, M., Martín, J., 2009. Immune activation affects chemical sexual ornaments of male Iberian wall lizards. *Naturwissenschaften* 96, 65–69.
- Louw, S., Burger, B.V., Le Roux, M., Van Wyk, J.H., 2007. Lizard epidermal gland secretions I: chemical characterization of the femoral gland secretion of the sun-gazer, *Cordylus giganteus*. *J. Chem. Ecol.* 33, 1806–1818.
- Martín, J., López, P., 2000. Chemoreception, symmetry and mate choice in lizards. *Proc. R. Soc. Lond. B* 267, 1265–1269.
- Martín, J., López, P., 2006a. Links between male quality, male chemical signals, and female mate choice in Iberian rock lizards. *Funct. Ecol.* 20, 1087–1096.

- Martín, J., López, P., 2006b. Vitamin D supplementation increases the attractiveness of males' scent for female Iberian rock lizards. Proc. R. Soc. Lond. B 273, 2619–2624.
- Martín, J., López, P., 2006c. Interpopulational differences in chemical composition and chemosensory recognition of femoral gland secretions of male lizards *Podarcis hispanica*: implications for sexual isolation in a species complex. Chemoecology 16, 31–38.
- Martín, J., López, P., 2006d. Age-related variation in lipophilic chemical compounds from femoral gland secretions of male lizards *Psammodromus algirus*. Biochem. Syst. Ecol. 34, 691–697.
- Martín, J., López, P., 2007. Scent may signal fighting ability in male Iberian rock lizards. Biol. Lett. 3, 125–127.
- Martín, J., López, P., 2010. Multimodal sexual signals in male ocellated lizards *Lacerta lepida*: vitamin E in scent and green coloration may signal male quality in different sensory channels. Naturwissenschaften 97, 545–553.
- Martín, J., López, P., 2011. Pheromones and reproduction in Reptiles. In: Norris, D.O., Lopez, K.H. (Eds.), Hormones and Reproduction of Vertebrates, Reptiles, vol. 3. Academic Press, San Diego, California, pp. 141–167.
- Martín, J., López, P., 2012. Supplementation of male pheromone on rock substrates attracts female rock lizards to the territories of males: a field experiment. PLoS One 7, e30108.
- Martín, J., López, P., 2013. Effects of global warming on sensory ecology of rock lizards: increased temperatures alter the efficacy of sexual chemical signals. Funct. Ecol. 27, 1332–1340.
- Martín, J., López, P., 2014. Pheromones and chemical communication in lizards. In: Rheubert, J.L., Siegel, D.S., Trauth, S.E. (Eds.), The Reproductive Biology and Phylogeny of Lizards and Tuatara. CRC Press, Boca Raton, Florida, USA, pp. 43–75.
- Martín, J., López, P., 2015. Condition-dependent chemosignals in reproductive behavior of lizards. Horm. Behav. 68, 14–24.
- Martín, J., Civantos, E., Amo, L., López, P., 2007a. Chemical ornaments of male lizards *Psammodromus algirus* may reveal their parasite load and health state to females. Behav. Ecol. Sociobiol. 62, 173–179.
- Martín, J., Moreira, P.L., López, P., 2007b. Status-signalling chemical badges in male Iberian rock lizards. Funct. Ecol. 21, 568–576.
- Martín, J., Chamut, S., Manes, M.E., López, P., 2011. Chemical constituents of the femoral gland secretions of male tegu lizards (*Tupinambis merianae*) (fam. Teiidae). Z. Naturforsch. C 66, 434–440.
- Martín, J., Castilla, A.M., Lopez, P., Al Jaidah, M., Mohtar, R., 2012. Lipophilic compounds in femoral gland secretions of spiny-tailed lizard, dhuh, *Uromastyx aegyptia microlepis* (Reptilia, Agamidae) from the Qatar desert. Qatar Found. Annu. Res. For 2012, EEP53.
- Martín, J., López, P., Garrido, M., Pérez-Cembranos, A., Pérez-Mellado, V., 2013a. Inter-island variation in femoral secretions of the Balearic lizard, *Podarcis lilfordi* (Lacertidae). Biochem. Syst. Ecol. 50, 121–128.
- Martín, J., Ortega, J., López, P., 2013b. Lipophilic compounds in femoral secretions of male collared lizards, *Crotaphytus bicinctores* (Iguania, Crotaphytidae). Biochem. Syst. Ecol. 47, 5–10.
- Martín, J., Ortega, J., López, P., 2013c. Chemical compounds from the preanal gland secretions of the male tree agama (*Acanthocercus atricollis*) (fam. Agamidae). Z. Naturforsch. C 68, 253–258.
- Martínez-Silvestre, A., Orós, J., 2002. Primera descripción de hiperplasia gingival en un lagarto gigante de El Hierro (*Gallotia simonyi*). Clin. Vet. Peq. Anim. 22, 170.
- Martínez-Silvestre, A., Rodriguez-Dominguez, M.A., Mateo, J.A., Pastor, J., Marco, I., Lavín, S., Cuenca, R., 2004a. Comparative haematology and blood chemistry of endangered lizards (*Gallotia* species) in the Canary Islands. Vet. Rec. 155, 266–269.
- Martínez-Silvestre, A., Silveira, L., Mateo, J.A., Urioste, J., Rodriguez-Dominguez, M.A., Pether, J., 2004b. Microbiología cloacal en lagartos gigantes amenazados de las Islas Canarias (Genero *Gallotia*) en cautividad. Rev. Esp. Herp 17, 29–37.
- Martínez-Silvestre, A., Marco, I., Rodriguez-Dominguez, M.A., Lavín, S., Cuenca, R., 2005. Morphology, cytochemical staining, and ultrastructural characteristics of the blood cells of the giant lizard of El Hierro (*Gallotia simonyi*). Res. Vet. Sci. 78, 127–134.
- Mason, R.T., 1992. Reptilian pheromones. In: Gans, C., Crews, D. (Eds.), Biology of the Reptilia, vol. 18. University of Chicago Press, Chicago, pp. 114–228.
- Mason, R.T., Parker, M.R., 2010. Social behavior and pheromonal communication in reptiles. J. Comp. Physiol. A 196, 729–749.
- Mayer, W., Pavlicev, M., 2007. The phylogeny of the family Lacertidae (Reptilia) based on nuclear DNA sequences: convergent adaptations to arid habitats within the subfamily Eremiinae. Mol. Phyl. Evol. 44, 1155–1163.
- McArdle, B.H., Anderson, M.J., 2001. Fitting multivariate models to community data: a comment on distance-based redundancy analysis. Ecology 82, 290–297.
- Müller-Schwarze, D., 2006. Chemical Ecology of Vertebrates. Cambridge University Press, Cambridge.
- Olsson, M., Madsen, T., Nordby, J., Wapstra, E., Ujvari, B., Wittsell, H., 2003. Major histocompatibility complex and mate choice in sand lizards. Proc. R. Soc. Lond. B (Suppl.) 270, 254–256.
- Pérez-Mellado, V., Romero-Benítez, M., Ortega, F., Martín-García, S., Perera, A., López-Vicente, M., Galache, C., 1999. El uso de los recursos tróficos en *Gallotia simonyi* (Sauria, Lacertidae) de la isla de El Hierro (Islas Canarias). In: López Jurado, L.F., Mateo, J.A. (Eds.), El Lagarto Gigante de El Hierro. Bases para su Conservación. Monografías de Herpetología. Asociación Herpetológica Española, Madrid, Spain, pp. 63–83.
- Pleguezuelos, J.M., Márquez, R., Lizana, M., 2002. Atlas y Libro Rojo de los Anfibios y Reptiles de España. Dirección General de Conservación de la Naturaleza. Asociación Herpetológica Española, Madrid.
- Roca, V., Orrit, N., Llorente, G.A., 1999. Parasitofauna del lagarto gigante de El Hierro, *Gallotia simonyi*. In: López Jurado, L.F., Mateo, J.A. (Eds.), El Lagarto Gigante de El Hierro. Bases para su Conservación. Monografías de Herpetología. Asociación Herpetológica Española, Madrid, Spain, pp. 127–139.
- Rodríguez-Domínguez, M.A., Molina-Borja, M., 1998. Reproduction of the endangered Hierro giant lizard *Gallotia simonyi machadoi*. J. Herp 32, 498–504.
- Rodríguez-Domínguez, M.A., Castillo, C., Coello, J.J., Molina-Borja, M., 1998. Morphological variation in the lacertid *Gallotia simonyi machadoi* and a comparison with the extinct *Gallotia simonyi simonyi* from El Hierro (Canary Islands). Herp J. 8, 85–91.
- Rodríguez-Domínguez, M.A., Trujillo González, D., Bohórquez-Alonso, M.L., Molina-Borja, M., 2014. Die aktuelle situation und die zukunftsperspektiven der rieseneidechsen von El Hierro und La Gomera, Kanarische Inseln. Teil 1: die El-Hierro-Rieseneidechse. Die Eidechse 25, 1–13.
- Weldon, P.J., Dunn, B.S., McDaniel, C.A., Werner, D.I., 1990. Lipids in the femoral gland secretions of the green iguana (*Iguana iguana*). Comp. Biochem. Physiol. B 95, 541–543.
- Weldon, P.J., Flachsbarth, B., Schulz, S., 2008. Natural products from the integument of nonavian reptiles. Nat. Prod. Rep. 25, 738–756.
- Wyatt, T.D., 2014. Pheromones and Animal Behaviour. Chemical Signals and Signatures Cambridge University Press, Cambridge.